

Novel Phase-Transition Behavior in an Aqueous Electrolyte Solution¹

J. Jacob^{2,3}, M. A. Anisimov^{2,4}, A. Kumar³, V. A. Agayan², and J. V. Sengers²

¹ Paper presented at the Fourteenth Symposium on Thermophysical Properties, June 25-30, 2000, Boulder, Colorado, U.S.A.

² Institute for Physical Science and Technology and Department of Chemical Engineering, University of Maryland, College Park, MD 20742, U.S.A.

³ Department of Physics, Indian Institute of Science, Bangalore 560 012, INDIA.

⁴ To whom correspondence should be addressed

ABSTRACT

We have investigated the near-critical behavior of the susceptibility of a ternary liquid mixture of 3-methylpyridine, water, and sodium bromide as a function of the salt concentration. The susceptibility was determined from light-scattering measurements performed at a scattering angle of 90° in the one-phase region near the locus of lower consolute points. A sharp crossover from asymptotic Ising behavior to mean-field behavior has been observed at concentrations ranging from 8 to 16.5 mass % NaBr. The range of asymptotic Ising behavior shrinks with increasing salt concentration and vanishes at a NaBr concentration of about 17 mass %, where completely mean-field-like behavior of the susceptibility is observed. A simultaneous pronounced increase in the background scattering at concentrations above 15 mass %, as well as a dip in the critical locus at 17 mass % NaBr, suggests that this phenomenon can be interpreted as mean-field tricritical behavior associated with the formation of a microheterogeneous phase due to clustering of the molecules and ions. An analogy with tricritical behavior observed in polymer solutions as well as a possibility of a charge-density-wave phase is also discussed. In addition, we, have observed a third soap-like phase on the liquid-liquid interface in several binary and ternary liquid mixtures.

KEY WORDS: aqueous ionic solutions; critical phenomena; crossover; light scattering.

1. INTRODUCTION

Critical phenomena in complex fluids such as ionic solutions, polymer solutions, micellar solutions, etc., continue to be an active topic of both theoretical and experimental research [1-6]. Asymptotically close to the critical point, fluids, both simple and complex are expected to belong to the Ising universality class [1,2]. In complex fluids, however, the interplay between universality caused by long-range fluctuations of the order parameter and a competing mesoscopic supra- or macromolecular structure may affect the approach to universal critical behavior. Hence, even within the Ising universality class, complex fluids may exhibit different crossover behavior upon approaching the critical point.

In polymer solutions the crossover from mean-field like behavior to asymptotic Ising behavior occurs closer to the critical point as the molecular weight of the polymer increases [7]. This crossover can be explained in terms of a competition between the correlation length ξ of the critical fluctuations and the radius of gyration (R_g) of the polymer molecules [7]. Polymer solutions exhibit a tricritical *theta* point in the limit of infinite molecular weight ($R_g \rightarrow \infty$) of the polymer [8]. The critical behavior near a tricritical point in three dimensions is mean-field one with small logarithmic corrections [9]. Hence the crossover observed in polymer solutions is, in fact, from the asymptotic Ising regime to the mean-field tricritical regime.

In ionic solutions the phase separation can be driven by either Coulombic interactions (low-dielectric-constant solvents) or solvophobic interactions (high-dielectric-constant solvents). In both cases, asymptotically close to the critical point, Ising-like behavior is expected due to a screening effect in the first case and due to short-range forces in the

second case [2]. Experiments on ionic solutions have revealed either Ising, mean-field or a crossover behavior between these two regimes [5]. A crossover from Ising behavior to mean-field behavior has been observed in several low-dielectric-constant “Coulombic” ionic solutions [10], although the correlation between the value of the dielectric constant and the crossover behavior has not been confirmed by other investigators.

To investigate the nature of criticality in aqueous ionic solutions, we have performed a series of light-scattering measurements in a ternary liquid mixture 3-methylpyridine (3MP) + water (H_2O) + sodium bromide (NaBr) [11]. The phase separation in this system is driven by solvophobic interactions in which hydrogen bonds between 3MP and H_2O play an important role. Light-scattering measurements reveal that the effective critical exponent γ_{eff} for the osmotic susceptibility χ exhibits a crossover from the 3-dimensional Ising asymptotic value ($\gamma = 1.24$) close to the critical temperature (T_c) to the mean-field value ($\gamma = 1.00$); this crossover behavior becomes more and more pronounced as the concentration of NaBr increases [11]. In this paper we show that for a sample with concentration of 17 mass % NaBr, complete mean-field behavior has been observed in the entire temperature range investigated. Moreover, a simultaneous pronounced increase of the background scattering appearing at the higher salt concentrations suggests that this phenomenon is to be interpreted as mean-field tricritical behavior associated with the appearance of a microheterogeneous phase due to clustering of ions and molecules.

2. EXPERIMENT

3-Methylpyridine and water are miscible in all proportions at atmospheric pressure, over the entire liquid-state temperature range (from freezing to boiling points of the liquids). However, a closed-loop immiscibility region with loop-size $\Delta T = T_U - T_L$ (T_U

and T_L being the upper and lower critical solution temperatures, respectively), appears when the mixture with 30 mass % 3MP is subjected to a pressure of approximately 1300 bar [12]. A similar effect can be realized by addition of ~ 0.1 mass percent of sodium chloride or ~ 0.4 mass percent of NaBr to the mixture (Fig. 1). With further addition of electrolyte, the gap ΔT increases. At the “double” critical point, at which the lower and upper critical points coincide, the asymptotic critical exponents are doubled [13]. Hence the effective values of the critical exponents will appear to be larger than their actual asymptotic values if ΔT is small. It has been demonstrated in earlier investigations of the system 3MP + H₂O + D₂O [13] that the critical exponents do not exhibit any noticeable increase beyond their Ising values when the loop-size (ΔT) is large (in practice, for $\Delta T \gtrsim 70 - 80^\circ\text{C}$). Hence, for large enough values of ΔT , the presence of the upper critical point can be neglected for measurements near the lower critical point. In the case of 3MP + H₂O + NaBr, it can be noted from Fig. 1 that even at a concentration of 5 mass % NaBr the size of the closed-loop is larger than 100°C . We also note that the lower critical temperatures are easily accessible experimentally over a wide range of electrolyte concentrations if NaBr is used in comparison to other electrolytes like sodium chloride, potassium bromide, sodium iodide, etc. The critical temperatures were also found to be stable in this system: the drift in T_L was less than 0.01°C over a period of two years.

Samples for our light-scattering measurements were prepared with 3MP (Aldrich, 99% pure), water (triple distilled in an all-quartz distiller) and analytical grade NaBr (99% pure). The critical composition of 3MP, $x_{\text{MP}} = (x_{\text{MP}})_c$, (x_{MP} being the mole fraction of 3MP in the ternary mixture) at each concentration of NaBr was accurately determined by choosing the concentration of the sample at which equal-volume phase separation occurs

at the phase-separation temperature. This procedure was repeated at each concentration of NaBr, since x_{MP} varies with the NaBr concentration. The lower critical temperatures were measured in a well-stirred liquid-paraffin thermostat with temperature stability better than $\pm 5\text{mK}$. The critical temperature was determined by the visual observation of the onset of phase separation and the eventual formation of an interface when the temperature is changed in steps of 2-3 mK. We have also ascertained the value of T_L before and after measuring the scattered-light intensity by monitoring the vanishing of the transmitted laser beam. For samples with $X > 0.16$ (X denotes the overall mass fraction of NaBr in the mixture) critical opalescence could be observed only in a narrow range of temperature close to the critical point, whereas for samples with $X < 0.16$ critical opalescence starts farther away from T_L .

The samples (volume $\sim 5\text{cc}$) were initially prepared in cylindrical pyrex glass cells and then transferred into the optical cells (volume $\sim 0.3\text{cc}$) by means of air-tight (Hamilton) syringes fitted with Millipore filters (pore size $0.2\ \mu\text{m}$). These cells were flame-sealed after the samples had been frozen in liquid nitrogen. A brass-block thermostat with a temperature stability better than $\pm 1\text{mK}$ was used in the light-scattering measurements. The laser beam ($\lambda = 632.8\ \text{nm}$) from a He-Ne laser (5 mW) was focused at the center of the sample. The intensity of scattered light (I) was measured at 90° in a photon-counting regime. The transmitted and incident beam intensities were measured with photodiodes. The light-scattering intensity data were normalized with respect to the incident-light intensity to account for fluctuations in the incident light. Corrections due to increased turbidity near the critical point have also been incorporated [14]. Figure 2 shows the total scattered intensity (I) as a function of $\tau = (T_L - T)/T$ for different X .

A crucial difference in the procedure of data acquisition of samples with $X \leq 0.16$ and $X > 0.16$ should be mentioned. Samples with $X \leq 0.16$ were stirred well for about 15-20 minutes in an ultrasonic agitator after preparation and also before starting each light-scattering run. However, in the case of samples with $X > 16$ mass % NaBr, a different procedure had to be adopted for measuring the scattered intensity. The samples were agitated in the ultrasonic agitator and then mounted in the thermostat at the temperature where the light-scattering measurements would begin. However, the samples were left undisturbed for about 24 hours before actually starting the measurements. Reproducible data from different experiments on the same sample were obtained only if this protocol was followed. Thermal equilibrium was indicated by the invariance of the scattered and transmitted intensities with time. The typical equilibration time for a temperature step of 0.1K (for the undisturbed sample) was ~ 20 -25 min which we found to be approximately the same for all concentrations. However, for samples with $X \geq 0.18$, it was not possible to obtain reproducible data even by following this procedure. This might be due to unusually large equilibration times for these samples. Hence, we have not attempted to analyze the data comprehensively for samples with $X \geq 0.18$.

3. RESULTS AND DISCUSSION

The intensity of light scattered by critical fluctuations (I_c) in fluids can be represented by

$$I_c = \frac{C\chi}{1 + q^2\xi^2} \quad (1)$$

where χ is the susceptibility, ξ the correlation length, C an experimental constant, and q the wave number, which is related to the scattering angle θ ($\theta = 90^\circ$ in our case) as $q =$

$4\pi n/\lambda_0 \sin(\theta/2)$, where $\lambda_0 = 632.8$ nm, the wavelength of the incident light (in vacuum), and where n is the refractive index. The dependence of the light-scattering intensity on the wave number q given by Eq. (1) is not exact [15] for large values of $q\xi$ but the difference is negligible for the wave numbers probed in our experiment. The intensity of light-scattering due to the critical fluctuations in the limit of zero wave number is proportional to the critical susceptibility χ , which diverges at the critical point. In a binary liquid solution the susceptibility is proportional to the osmotic compressibility

$$\left(\frac{\partial x_2}{\partial \mu_{2-1}} \right)_{P,T}$$

where x_2 is the concentration (mole fraction) of the solute and $\mu_{2-1} = \mu_2 - \mu_1$,

the difference of solute/solvent chemical potentials conjugate to x_2 . In a ternary liquid solution, the critical susceptibility is proportional to a generalized osmotic

$$\text{compressibility defined as } \chi = \left(\frac{\partial x_2}{\partial \mu_{2-1}} \right)_{P,T,\mu_{3-1}}$$

with $\mu_{3-1} = \mu_3 - \mu_1$ being the chemical

potential difference of the third component and the solvent conjugate to x_3 (since $d\mu_1 + x_2 d\mu_{2-1} + x_3 d\mu_{3-1} = 0$ at constant P and T). The generalized susceptibility and the

correlation length exhibit the asymptotic singular behavior $\chi = \Gamma_0 \tau^{-\gamma}$ and $\xi = \xi_0 \tau^{-\nu}$ respectively, where Γ_0 and ξ_0 are system-dependent critical amplitudes, and where the

critical exponents $\gamma = 1.24$ and $\nu = 0.63$ are universal, provided that the chemical potential difference $\mu_{3-1} = \mu_3 - \mu_1$ is constant. However, the light-scattering measurements

are performed at constant overall concentration ($x_3 = \tilde{X}$, the overall mole fraction of NaBr) and, as a consequence, the experimentally observed asymptotic critical-exponent

values are affected by Fisher renormalization [16]. As a consequence the susceptibility exponent changes its asymptotic value from γ to $\gamma(1-\alpha)$ with $\alpha = 0.11$, the critical

exponent of the so-called weakly divergent susceptibility [17]. In our data analysis, we have accounted for this Fisher renormalization by expressing the experimental temperature scale $\tau(\tilde{X})$ through the theoretical temperature scale $\tau = \tau(\mu_{3-1})$ [18]:

$$\frac{\tau(\tilde{X})}{\tau_2} \cong \left(\frac{\tau(\mu_{3-1})}{\tau_2} \right)^{1-\alpha} \left[1 + \left(\frac{\tau(\mu_{3-1})}{\tau_2} \right)^\alpha \right] \quad (2)$$

where the characteristic temperature τ_2 is defined by

$$\tau_2 = \left[A_0 \tilde{X} \left(1 - \tilde{X} \left(\frac{1}{T_c} \frac{dT_c}{d\tilde{X}} \right)^2 \right)^{1/\alpha} \right] \quad (3)$$

where A_0 is the critical amplitude of the weakly-divergent ($\sim \tau^\alpha$) susceptibility which is related to the amplitude ξ_0 of the power law for the correlation length by a universal two-scale-factor relation [11]. The extent and observability of Fisher renormalization depends on the parameter τ_2 , and, hence, on the concentration \tilde{X} and the slope of the line of critical temperatures versus \tilde{X} , $(dT_c/d\tilde{X})$. In the mean-field regime where $\alpha = 0$, Fisher renormalization vanishes.

In analyzing data in terms of Eq. (1), χ and ξ have been represented by equations from a crossover theory, which recovers Ising asymptotic behavior with Wegner corrections close to the critical point and mean-field behavior far away from the critical point [19]. For samples with $X \geq 0.16$, the total intensity includes a contribution from background scattering also. The intensity of the background scattering (I_0) was calculated as the difference between the total intensity (I) and the intensity (I_c) from the critical

fluctuations. As an alternative approach, we also fitted the data, including those far away from the critical point, to the expression $I = I_c + I_0$ where I_0 was represented by

$$I_0 = m_1(\tau - \tau_0) + m_2(\tau - \tau_0)^2 \quad (4)$$

where τ_0 , m_1 , and m_2 are adjustable parameters (Table I). Physically, Eq. (4), which is similar to the virial-like expression expected near the critical micellar concentration [20], corresponds to the assumption that the formation of “clusters” emerges at a certain $\tau = \tau_0(X)$. The crossover behavior of χ and ξ obtained with these two alternative treatments of the background scattering appeared to be essentially the same. From Figs. 2 and 3, it can be seen that I_0 is negligible at $X \leq 0.15$, but it becomes increasingly prominent with increase in X . Small-Angle X-ray Scattering (SAXS) studies in these solutions have provided evidence for cluster formation in samples with $X \geq 0.12$ [21]. A systematic increase in cluster size with concentration was shown in the range $0.12 \leq X \leq 0.17$. This clustering was explained as due to the formation of 3MP-rich clusters resulting from a preferential solvation of Na^+ and Br^- ions by water molecules [21]. Evidence of clustering induced by potassium chloride in a somewhat similar solution of 1-propanol and water has been provided by small angle neutron scattering measurements also [22]. Therefore, the temperature $\tau = \tau_0(X)$ in Eq. (4) can be interpreted as the temperature of cluster formation and can be determined experimentally. Figure 3 shows the relative excess scattering intensity, defined as $I_0/(I_c+I_0)$, as a function of the distance to the critical temperature. It can be seen that the emergence of the “background” scattering coincides with the critical temperature somewhere above $X = 0.165$ and slightly below $X = 0.17$.

The crossover to mean-field behavior is generally controlled by two physical parameters, namely a rescaled coupling constant \bar{u} which reflects the range of molecular

forces and a so-called “cutoff” [4,7,19]. In simple fluids the “cutoff” is related to the molecular size and the trend to mean-field critical behavior is controlled by the coupling constant only. In complex fluids the “cutoff” is associated with the characteristic supramolecular or macromolecular length $\xi_D = v_0^{1/3} \Lambda^{-1}$ representing a crossover length scale [1], with v_0 being an average molecular volume ($v_0^{1/3} \cong 3.5 \text{ \AA}$ for our samples [11]) and Λ is a dimensionless characteristic cutoff wave-number. If ξ_D is large enough, it competes with the correlation length of the critical fluctuations and controls a crossover to mean-field tricritical (or, more generally, multicritical) behavior. When both ξ and ξ_D diverge, one will recover a mean-field tricritical point. It is important to emphasize the physical difference between approaching mean-field behavior due to the long-range nature of the molecular forces (small \bar{u} while $\bar{\xi}_0/\xi_D$ is not small) and that due to a small ratio ξ_0/ξ_D (i.e. the ratio $\xi_D/\bar{\xi}_0$ is large, while \bar{u} is not necessarily small). The former case is the conventional crossover to mean-field critical behavior. Complete crossover of such a kind has been observed in the three-dimensional Ising model with variety of interaction ranges [23,24]. The latter case of large and eventually diverging ξ_D corresponds to approaching a special kind of mean-field behavior, namely tricritical mean-field behavior. A typical and conceptually well-understood example is the crossover to theta-point tricriticality in polymer solutions [25]. One may obtain an empirical crossover temperature τ_\times as the value of τ where the effective susceptibility exponent, defined as $\gamma_{\text{eff}} = -\tau d \ln \chi/d\tau$, and recovered after accounting for Fisher renormalization, exhibits an inflection point as a function of τ [7,11]. We have found crossover behavior from Ising critical behavior asymptotically close to the critical temperature to mean-field critical behavior further away from the critical temperature at

all salt concentrations up to and including $X = 0.165$. However, the range of Ising critical behavior shrinks with increasing salt concentration. In Fig. 4 we compare the deviations from Ising behavior at the higher salt concentrations with those at the lowest salt concentration $X = 0.08$. At $X = 0.08$, all experimental data are in the crossover regime, at $X = 0.16$ a substantial fraction of the experimental data exhibits mean-field critical behavior, at $X = 0.165$ the experimental data follow the mean-field curve except for the two data points closest to T_c , and at $X = 0.17$ the parameter Λ becomes insignificantly small and the crossover temperature becomes undetectable (Table I). Thus within our experimental resolution, complete mean-field critical behavior is observed at $X = 0.17$. In Fig. 5 we plot the effective susceptibility exponent γ_{eff} as a function of the reduced temperature τ . It is evident from this figure that the crossover from Ising-like to mean-field behavior becomes more pronounced at larger NaBr concentrations. For the samples with $X = 0.16$, and 0.165 , the crossover is practically completed within the critical domain. At $X = 0.17$ we did not find any trend to Ising behavior near the critical point and the observed susceptibility retained mean-field behavior upon the approach to T_c . In addition the crossover parameter Λ becomes insignificantly small and the crossover temperature τ_x becomes undetectably small. In Fig. 6 we show the crossover temperature τ_x and the crossover length scale ξ_D as a function of the salt concentration. As follows from the theory of crossover from Ising critical behavior to mean-field tricritical behavior [4], at $\tau = \tau_x$ the correlation length ξ of the critical fluctuations is equal to the competing correlation length ξ_D associated with an additional order parameter. It is seen that the crossover temperature τ_x vanishes and the crossover length ξ_D diverges at a salt

concentration between 16.5 and 17%. Thus both ξ and ξ_D diverge at this point, giving strong evidence for tricriticality.

The most interesting and intriguing question to be discussed is the physical nature of the discovered mean-field like multicritical point. There are several possible scenarios of multicritical behavior in a ternary mixture, each of them requiring further investigation. Here we just briefly discuss some phenomenological consequences of these scenarios.

- (a) A fluid-mixture vapor-liquid-liquid tricritical point. In a ternary fluid mixture there might exist a tricritical point, defined as a point at which two lines of critical end points merge and three near-critical fluid phases become identical [26]. Such a scenario does not look realistic for a ternary liquid solution at atmospheric pressure, as one of the fluid phases along the line of critical end points should originate from the near-critical vapor phase.
- (b) A tricritical point caused by a coupling between two order parameters belonging to different universality classes. If we assume, that an additional (vector-like) order parameter emerges in the system, there might be a coupling between the scalar order parameter (concentration) associated with the liquid-liquid phase separation and this vector-like order parameter associated with some kind of structural ordering in the one-phase region. A projection of this hypothetical phase diagram on $T - \mu_{3-1}$ plane is shown in Fig. 7. In this figure the λ -line (dotted line) is a line of second order phase-transitions between two macroscopically homogeneous, but structurally different phases. A coupling between formation of a microheterogeneous (e.g., micellar-like or microemulsion sponge-like) phase and liquid-liquid phase separation can lead to tricritical behavior [20]. At the tricritical point the λ -line merges with two wings each

of them being the critical liquid-liquid phase-separation line. Each wing belongs to an opposite sign of the “ordering field” h , a linear combination of μ_{3-1} , μ_{2-1} , and T . At negative h , two molecular solutions coexist in the two-phase region. At positive h , two microheterogeneous solutions coexist. Along the λ -line, the “ordering field” h is zero with the tricritical point separating the second-order and first-order transitions (shown in Fig. 7 as dashed line). Along the first-order transition line the molecular solution and microheterogeneous solutions are in liquid-liquid coexistence and change in the fluid structure is accompanied by phase separation. Such a diagram can be compared with the actual phase diagram of the ternary mixture 3MP + H₂O + NaBr shown in Fig. 1.

- (c) If the λ -line shown in Fig. 7 corresponds to a transition between a molecular-solution phase and a microheterogeneous phase characterized by an equilibrium mesoscopic wave number, the tricritical point may become a so-called Lifshitz point [27]. In this particular case, the right wing of the critical phase separation line shown in Fig. 7 will become a first-order liquid-liquid phase separation between the molecular solution and a microheterogeneous phase. The liquid-liquid coexistence between two microheterogeneous phases (the dotted curve in figure 7) will not appear. Physically, it means that the microheterogeneous structure becomes unstable with onset of phase separation. The first-order phase separation will become continuous at a Lifshitz point. Such a scenario seems to be quite plausible. The possibility of a multicritical Lifshitz point in which two fluid phases in an electrolyte solution coexist with a microheterogeneous charge-density wave phase has been suggested by Nabutovskii *et al.* [28] and discussed by Höye and Stell [29] and Fisher [1]. Evidence for tricritical

behavior in restricted primitive models for ionic fluids has been reported by Dickman and Stell [30] and by Panagiotopoulos and Kumar [31]. In our system at higher salt concentration the 3-methylpyridine molecules are assumed to be shielded from the ions by water molecules. In turn, the ions may form a double layer over the water molecules forming a charge layering or microemulsion-sponge-like domains.

4. FORMATION OF A THIRD PHASE

Another intriguing phenomena that we encountered during our investigations of samples with $X \geq 0.16$ was the observation of small amount of a third soap-like phase on the interface. This phase was first observed in the system 3-Methylpyridine + Water + Sodium Bromide at approximately 16.5 mass percent of NaBr, i.e., very close to the discovered multicritical point. It appeared on the interface immediately after phase separation of the sample. The experiments were performed near the lower consolute temperature. The samples were heated slowly from the one-phase region to the critical temperature. Though the third phase was observed on the interface very close to the critical temperature, the samples were overheated approximately 0.5K into the two-phase region in order to observe the third-phase clearly. The samples were filtered with 0.2 μm Millipore filters to ensure that dust was avoided in the samples. These samples were also prepared from chemicals of different purity: NaBr – 99.6%, and 99.99%, MP – 98%, 99%, and 99.7%. The third phase was observed irrespective of any of the above mentioned conditions. The amount of the third phase observed was investigated as a function of the concentration of NaBr. Samples with NaBr concentration up to 27 mass % have been investigated. It was observed that there is no appreciable increase in the amount of the third phase when the concentration of NaBr is increased beyond 17 mass

% . The third phase was also observed in samples with NaBr concentrations as low as 0.1 mass % though the amount of the third phase was appreciably lower. If the salt NaCl was used instead of NaBr, the third phase was observed in these samples as well. We also prepared samples with 3MP and heavy water (D_2O) where the phase separation occurs without salt and traces of the third phase were seen even in these samples. Under a magnification of 10 times, the third phase has a whitish fibrous appearance. If, after the third phase appears on the interface, the samples are cooled back into the one-phase region and shaken well, very fine fiber-like particles could be seen in the samples. We filtered these samples again and repeated the experiments. It was observed that the third phase forms again in these samples and there is no appreciable change in the value of T_c or the position of the interface. The samples were prepared in different kinds of glass tubes and also in quartz tubes and the third phase was observed to form irrespective of the kind of tube used.

Visual investigations were also carried out on different liquid mixtures in an attempt to characterize this solid-like phase. Samples of 2,6-lutidine + water were prepared to check whether the third phase is seen in this system, although a strong wetting phenomenon might prevent such an observation. The third phase was not observed in this sample. But, on addition of approximately 3 mass % NaBr to this binary mixture, the third phase clearly appeared at the interface. For a 10% solution of NaBr, the amount of the third phase seemed to be larger than in the case of a 3% solution. It may be that the amount of the third phase was so small in 2,6-lutidine + water, without any NaBr that it was not visually noticeable.

The third phase has also been observed in mixtures of isobutyric acid (IBA) + water. These samples were prepared with IBA (99%, Aldrich) and deionized water obtained from an Alpha-Q water purification system. The samples were also filtered using 0.2 μ m filters. The phenomenon was checked in both pyrex and quartz cells and no difference has been observed. We have also reexamined two samples of isobutyric acid + water and isobutyric acid + heavy water studied by Greer many years ago [32]. These samples had been prepared in pyrex glass tubes and they had not been filtered. Upon close examination, traces of a third phase were observed in these samples also.

The third phase has not been observed in solutions of 3-methylpentane + nitroethane. Since the refractive index of these two liquids is very close to each other, critical opalescence is greatly reduced and it was possible to observe the interface at temperatures very close to T_c . But we did not detect any traces of a soap-like phase at the interface in these solutions. It is not clear yet whether the formation of a soap-like phase at the interface is associated with the formation of a microheterogeneous phase in the bulk observed in the system 3MP + H₂O + NaBr and whether it is related to wetting phenomena.

ACKNOWLEDGMENTS

We are indebted to R. W. Gammon for providing us with the capability of observing phase separation in our liquid mixtures, to S. C. Greer for making her samples of isobutyric acid + water available, and to M. E. Fisher, J. M. H. Levelt Sengers, A. Oleinikova, R. A. Narayanan, G. Stell, and H. Weingärtner for stimulating discussions. The research at the University of Maryland is supported by the National Science Foundation under Grant No. CHE-9805260. The research at the Indian Institute of

Science has been supported by the Department of Science and Technology, Government of India.

REFERENCES

1. M. E. Fisher, *J. Stat. Phys.* **75**, 1 (1994).
2. G. Stell, *Phys. Rev. A* **45**, 7628 (1992); *J. Stat. Phys.* **78**, 197 (1995).
3. H. Weingärtner, M. Kleemeier, S. Wiegand, W. Schröer, *J. Stat. Phys.* **78**, 169 (1995).
4. M. A. Anisimov, A. A. Povodyrev, and J. V. Sengers, *Fluid Phase Equilibria* **158-160**, 537 (1999); M. A. Anisimov, *J. Phys.: Cond. Matt.* **12**, A451 (2000).
5. J. M. H. Levelt Sengers, A. H. Harvey, and S. Wiegand, in *Equations of State for Fluids and Fluid Mixtures*, J. V. Sengers, R. F. Kayser, C. J. Peters, H. J. White Jr., eds. (Elsevier, Amsterdam, in press).
6. R. G. Rubio and F. Ortega, *J. Phys.: Cond. Matt.* **12**, A459 (2000).
7. Y. B. Melnichenko, M. A. Anisimov, A. A. Povodyrev, G. D. Wignall, J. V. Sengers, and W. A. Van Hook, *Phys. Rev. Lett.* **79**, 5266 (1997).
8. P. G. de Gennes, *Scaling Concepts in Polymer Physics* (Cornell University Press, Ithaca NY, 1979).
9. I. D. Lawrie and S. Sarbach, in *Phase Transitions and Critical Phenomena*, C. Domb and J. L. Lebowitz, eds. (Academic Press, New York, 1984), vol. 9, p.2.
10. T. Narayanan and K. S. Pitzer, *Phys. Rev. Lett.* **73**, 3002 (1994); *J. Chem. Phys.* **102**, 8118 (1995).
11. J. Jacob, A. Kumar, M. A. Anisimov, A. A. Povodyrev, and J. V. Sengers, *Phys. Rev. E* **58**, 2188 (1998).
12. G. M. Schneider, *Ber. Bunsenges. Phys. Chem.* **76**, 325 (1972).
13. B. V. Prafulla, T. Narayanan, and A. Kumar, *Phys. Rev. A* **45**, 1266 (1992).

14. A. J. Bray and R. F. Chang, *Phys. Rev. A* **12**, 2594 (1975).
15. M. E. Fisher and R. J. Burford, *Phys. Rev.* **156**, 583 (1967).
16. M. E. Fisher, *Phys. Rev.* **176**, 237 (1968).
17. M. A. Anisimov, E. E. Gorodetskii, V. D. Kulikov, and J. V. Sengers, *Phys. Rev. E* **51**, 1199 (1995).
18. H. Y. Cheng, M. A. Anisimov, and J. V. Sengers, *Fluid Phase Equilibria* **128**, 67 (1997).
19. M. A. Anisimov, A. A. Povodyrev, V. D. Kulikov, and J. V. Sengers, *Phys. Rev. Lett.* **75**, 3146 (1995).
20. M. A. Anisimov, A. S. Kurlandsky, and N. F. Kazakova, *Mol. Cryst. Liq. Cryst.* **159**, 87 (1988).
21. J. Jacob, A. Kumar, S. Asokan, D. Sen, R. Chitra, and S. Mazumder, *Chem. Phys. Lett.* **304**, 180 (1999).
22. M. Misawa, K. Yoshida, K. Maruyama, H. Munemura, and Y. Hosokawa, *J. Phys. Chem. Solids* **60**, 1301 (1999).
23. E. Luijten and K. Binder, *Phys. Rev. E* **58**, R4060 (1988).
24. M. A. Anisimov, E. Luijten, V. A. Agayan, J. V. Sengers, and K. Binder, *Phys. Lett. A* **264**, 63 (1999).
25. A. A. Povodyrev, M. A. Anisimov, and J. V. Sengers, *Physica A* **264**, 345 (1999).
26. C. M. Knobler and R. L. Scott, in *Phase Transitions and Critical Phenomena*, C. Domb and J. L. Lebowitz, eds. (Academic Press, New York, 1984), vol. 9, p.164.
27. L. D. Landau and E. M. Lifshitz, *Statisticheskaya Fizika*, Part 1, 5th Russian Edition (Fizmatlit, Moscow, 1995).

28. V. M. Nabutovskii, N. A. Nemov, and Yu. G. Peisakhovich, *Zh. Eksp. Teor. Fiz.* **79**, 2196 (1980) [*Sov. Phys. JETP* **52**, 1111 (1980)]; *Phys. Lett. A* **79**, 98 (1980); *Mol. Phys.* **54**, 979 (1985).
29. J. S. Hoye and G. Stell, *J. Phys. Chem.* **94**, 7899 (1990).
30. R. Dickman and G. Stell, in *Simulation and Theory of Electrostatic Interactions in Solution*, L. R. Pratt and G. Hummer, eds. (American Institute of Physics, New York, 1999), p. 225; G. Stell, in *New Approaches to Problems in Liquid State Theory*, C. Caccamo, J. P. Hansen, and G. Stell, eds. (Kluwer, Dordrecht, 1999), p. 71.
31. A. Z. Panagiotopoulos and S. K. Kumar, *Phys. Rev. Lett.* **83**, 2981 (1999).
32. S. C. Greer, *Phys. Rev. A* **14**, 1770 (1976).

Table I. Parameters for description of crossover behavior of the light-scattering intensity in 3-methylpyridine-water-NaBr solutions.

Critical Parameters		Crossover Parameters			Background light-scattering parameters		
X	$T_c(\text{K})$	τ_x	$\bar{\xi}_0 (\text{\AA})$	$\xi_D (\text{\AA})$	τ_0	m_1	m_2
0.16	306.435 ± 0.004	3×10^{-3}	4.60	70.3	2.1	0.14	0.006
0.165	305.131 ± 0.004	3×10^{-4}	5.25	230	0.96	0.17	0.016
0.17	304.628 ± 0.004	$< 3 \times 10^{-5}$	5.67	$> 3 \times 10^3$	-0.15	0.04	0.061
0.18	303.888 ± 0.004	not defined			-0.22	0.25	0.19

FIGURE CAPTIONS

Fig. 1. Phase diagram of the 3-methylpyridine + water + NaBr system. The insert shows an expanded view of the range of NaBr concentrations where light-scattering measurements have been performed. The solid curve indicates the critical phase-separation line and the dashed curve bounds the temperature and salt concentrations where a microheterogeneous phase exists. The crosses indicate the temperatures corresponding to $\tau = \tau_0$ according to Eq. (2) where clustering appears. The horizontal bar indicates the concentration range at $T = 298$ K, where clustering appears according to SAXS measurements.

Fig. 2. Light-scattering intensity (I) as a function of τ for samples with seven different mass fractions of NaBr. The symbols denote the experimental data and the curves represent values calculated from the crossover model with a correction for enhanced background scattering due to cluster formation for $X \geq 0.16$.

Fig. 3. The relative excess scattering $[I_0/(I_0 + I_c)]$ as a function of the distance to the critical point $(T-T_L)$ for samples with $X \geq 0.16$. The solid and dashed curves represent fits to the data in terms of Eq. (4).

Fig. 4. Deviations of the scattering intensity from asymptotic Ising critical behavior as a function of $\tau = (T_c - T)/T$ for four different salt concentrations. The symbols designate experimental data. The dashed curves represent mean-field behavior and the solid curve represents the actual behavior as calculated from the crossover model with a correction for enhanced background scattering for $X \geq 0.16$. The solid curve for $X = 0.17$ corresponds to a crossover fit with the insignificant crossover temperature $\tau_x = 3 \times 10^{-6}$ and coincides with the mean-field (dashed) curve.

Fig. 5. Effective susceptibility exponent $\gamma_{\text{eff}} = -\tau d \ln\chi/d\tau$ for seven different salt concentrations calculated from the crossover model after accounting for Fisher renormalization.

Fig. 6. The crossover temperature τ_x and the crossover length-scale ξ_D as a function of the salt concentration. The solid curve represents an empirical power law $\tau_x \propto (X_0 - X)^{0.8}$ with $X_0 = 0.1652$. The dashed curve represents a power law $\xi_D \propto (X_0 - X)^{-1/2}$.

Fig. 7. General schematic phase diagram representing a ternary system with a tricritical point (TCP) emerging as a result of coupling between two different order parameters.




Fig. 1




Fig. 2




Fig. 3







Fig. 4




Fig. 5




Fig. 6




Fig. 7